

Cat.No # MGF9012

Lot.No : Refer to Vial



Packaging: 10µg \ 50µg \ 500µg

Description

Human Vascular endothelial growth factor (VEGF), also known as VEGF-A and vascular permeability factor (VPF), belongs to the platelet-derived growth factor family of cysteine-knot growth factors. It is a potent activator in vasculogenesis and angiogenesis both physiologically and pathologically. VEGF-A has 8 differently spliced isoforms, of which VEGF165 is the most abundant one. VEGF165 is a disulfide-linked homodimer consisting of two glycosylated 165 amino acid polypeptide chains. VEGF stimulates the cellular response through binding to tyrosine kinase receptors VEGFR1 and VEGFR2 on the cell surface. It is widely accepted that VEGFR2 mediate almost all of the known cellular responses to VEGF while the function of VEGFR1 is less defined and is thought to modulate the VEGFR2 signaling.

Amino acid sequence

VEGF165 (Ala27-Arg191)

Origin

Recombinant Human Vascular Endothelial Growth Factor Isoform 165/VEGF165 was produced in human cells that were transfected with an expression plasmid encoding Human VEGF165 (Ala27-Arg191).

Quality control

Greater than 95% as determined by SEC-HPLC and reducing SDS-PAGE.

Dissolution

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Store condition

Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20°C for 3 months.

For Research Use Only. Not for use in diagnostic procedures.

